

OxThera AB						
Name of study	Objective	Major Inclusion criteria	Major exclusion criteria	Major time points	Centers participating in Europe	More information available at
OC5-DB-02 (ePHex study)	To evaluate the efficacy of Oxabact [®] following 52 weeks treatment in patients with maintained kidney function but below the lower limit of the normal range (eGFR < 90 ml/min/1.73 m ²) and a total plasma oxalate concentration ≥ 10 µmol/L at baseline.	<ol style="list-style-type: none"> 1. Signed informed consent 2. Diagnosis of PH. 3. eGFR < 90 ml/min/1.73 m². 4. Total plasma oxalate concentration ≥10 µmol/L 5. ≥ 2 years of age 6. stable vitamin B6 for at least 3 months prior to screening and must not change the dose during the study. 	<ol style="list-style-type: none"> 1. Inability to swallow size 4 capsules. 2. Previous transplantation (solid organ or bone marrow). 3. Patients requiring dialysis or at immediate risk for kidney failure 4. Secondary hyperoxaluria 5. Use of antibiotics to which O. formigenes is sensitive 	Baseline, week 24, week 52	<p>UK:</p> <ul style="list-style-type: none"> • Nottingham Children's Hospital • Royal Free Hospital -UCL <p>Germany: Kindernierenzentrum Bonn</p> <p>Belgium: Centre Hospitalier Universitaire de Liège</p> <p>USA: Vanderbilt University Hospital</p> <p>Spain: Hospital Universitari Vall d' Hebron</p>	www.oxthera.com www.clinicaltrials.gov

					Tunisia: <ul style="list-style-type: none"> • Hedi Chaker University Hospital, Sfax • Hôpital Charles-Nicolle, Tunis • Hôpital Sahloul, Sousse 	
OC5-OL-02 (ePHex-OLE study)	To evaluate the efficacy of Oxabact [®] following two years continued treatment in subjects who have completed the Oxabact [®] OC5-DB-02 (ePHex) study.	<ol style="list-style-type: none"> 1. Signed informed consent 2. Participation in and completion of study ePHex. 	Same as above	Baseline, week 52, week 104	Same as above	www.oxthera.com www.clinicaltrials.gov